FORT LEE SCHOOL DISTRICT FORT LEE, NEW JERSEY

REQUEST FOR GIVING MEDICATION AT SCHOOL FORM 02-D-18

This form is required for all over-the-counter and/or prescription medication(s) to be administered during school hours. The medication(s) will be supplied by the parents and brought to the school nurse in the <u>original</u> container appropriately labeled by the pharmacy and physician. All medication must be picked up at the end of the school year.

^		Name	School Doctor
Received	by school and reviewed by		School Nurse-teacher
Par	ent/ Guardian Signature	Emergency contact number	r Date
•	medication.	se/school physician perm	Initial: Initial: Initial:
	tion is to be completed by the Parcinitial the following:	ent/ Legal Guardian	
PHY	SICIAN PRINTED NAME		PHYSICIAN STAMP (TO INCLUDE ADDRESS & PHONE NUMBE
PHYS	SICIAN SIGNATURE	DATE	
Trips (including overnight trips)	☐ Other prescribed time	period:
Dates t	to be dispensed (Please che	ck): School year	to □ Half days □ Field
Interve	ention to be rendered for an	adverse reaction:	
Possib	le side effects of medicatio	n:	
Route:	Time to be a	administered:	am/pm (please circle)
Name	of Medication:	Dos	e to be administered:
Diagno	osis/Medical Condition:		
Allergi	les:	Grade:	Current Weight:
Studen	t's Name:		Date of Birth:

Data



FOOD ALLERGY & ANAPHYLAXIS EMERGENCY CARE PLAN

Other (e.g., inhaler-bronchodilator if wheezing):

F. OF ALLERS RESERVED & COLUMNIC				
Name:	D.O.B.: PLACE PICTURE			
Allergic to:				
Weight:Ibs. Asthma:				
NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.				
Extremely reactive to the following allergens:				
THEREFORE:				
☐ If checked, give epinephrine immediately if the allergen was LIKELY eaten, for ANY symptoms.				
☐ If checked, give epinephrine immediately if the allergen was DEFINITELY eaten, even if no symptoms are apparent.				
FOR ANY OF THE FOLLOWING:	MILD SYMPTOMS			
SEVERE SYMPTOMS				
LUNG HEART THROAT MOUTH	NOSE MOUTH SKIN GUT Itchy or Itchy mouth A few hives, Mild			
Shortness of Pale or bluish Tight or hoarse Significant	runny nose, mild itch nausea or			
breath, wheezing, skin, faintness, throat, trouble swelling of the repetitive cough weak pulse, breathing or tongue or lips	sneezing discomfort			
dizziness swallowing	FOR MILD SYMPTOMS FROM MORE THAN ONE SYSTEM AREA, GIVE EPINEPHRINE.			
OR A COMBINATION	FOR MILD SYMPTOMS FROM A SINGLE SYSTEM			
SKIN GUT OTHER of symptoms	AREA, FOLLOW THE DIRECTIONS BELOW:			
Many hives over Repetitive Feeling from different body, widespread vomiting, severe something bad is body areas.	Antihistamines may be given, if ordered by a healthcare provider.			
redness diarrhea about to happen, anxiety, confusion	2. Stay with the person; alert emergency contacts.			
0 0	3. Watch closely for changes. If symptoms worsen,			
1. INJECT EPINEPHRINE IMMEDIATELY.	give epinephrine.			
Call 911. Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders	MEDICATIONS/DOSES			
arrive. Consider giving additional medications following epinephrine:	Epinephrine Brand or Generic:			
» Antihistamine » Inhaler (bronchodilator) if wheezing	Epinephrine Dose: 0.1 mg IM 0.15 mg IM 0.3 mg IM			
Lay the person flat, raise legs and keep warm. If breathing is	Antihistamine Brand or Generic:			
difficult or they are vomiting, let them sit up or lie on their side. • If symptoms do not improve, or symptoms return, more doses of	Antihistamine Dose:			

epinephrine can be given about 5 minutes or more after the last dose.

Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

Alert emergency contacts.